

9.1. Exposure scenario 1: Widespread use by professional workers

Outdoor use – direct application of solid Mn fertilizers to soil; surface spreading

Product category used: PC 12: Fertilizers

Sector of use: SU 1: Agriculture, forestry, fishery

| Environment contributing scenario(s): | | |
|---------------------------------------|---|---|
| CS 1 | Outdoor use – direct application of solid Mn fertilizers to soil; surface spreading | ERC 8e Fertilizers Europe SPERC 8e.1.v2 |

9.1.1. Env CS 1: Outdoor use - direct application of solid Mn fertilizers to soil; surface spreading (Fertilizers Europe SPERC 8e.1.v2; ERC 8e)

9.1.1.1. Conditions of use

| | |
|--|--|
| Product (article) characteristics | <ul style="list-style-type: none"> Solid fertilizers intended for outdoor use (in a.o. agriculture, forestry, horticulture, gardens, golf courses) by consumers and professionals. Farmers are considered professional users. |
| Amount used, frequency and duration of use (or from service life) | <ul style="list-style-type: none"> Daily local widespread use amount: not relevant Substance use amount expressed as yearly fertilizer application rate (kg/ha/year): 3.0 kg Mn/ha/year Only to be applied if manganese deficiency is detected Number of emission days per year: 1-5 applications per year; depending on crop type and agricultural soil characteristics |
| Conditions and measures related to biological sewage treatment plant | <ul style="list-style-type: none"> Biological STP: None [Effectiveness Water: 0%] |
| Conditions and measures related to external treatment of waste (including article waste) | Particular considerations on the waste treatment operations: Other <ul style="list-style-type: none"> Dispose of waste product or used containers according to local regulations Service life is not applicable to fertilizers |
| Other conditions affecting environmental exposure | <ul style="list-style-type: none"> Outdoor use [OOC01] Controlled application to agricultural soil [OOC25] |

9.1.1.2. Releases

The local releases to the environment are reported in the following table.

Table 9.1.1 Local releases to the environment

| Release | Release estimation method | Explanations |
|-------------------|---------------------------|--|
| Water | Estimated release factor | Release factor before on site RMM: 0% Release factor after on site RMM: 0% Local release rate: not applicable |
| Air | Estimated release factor | Release factor before on site RMM: 0% Release factor after on site RMM: 0% |
| Agricultural soil | Estimated release factor | Release factor after on site RMM: 100% (direct application to agricultural soil, intentional release) |

9.1.1.3. Exposure and risks for the environment and man via the environment

The exposure concentrations and risk characterization ratios (RCR) are reported in the following table. The exposure estimates have been obtained with the Fertilizer Environmental Exposure (FEE) Tool v1.2 unless stated otherwise. The environmental risk assessment for manganese follows an added risk approach and therefore no natural Mn background concentration in soil, sediment or water is considered for exposure and risk assessment.

Table 9.1.2. Exposure concentrations and risks for the environment

| Protection target | Exposure concentration | Risk quantification |
|-------------------------|---|---------------------|
| Fresh water | Local PEC: 2.5 µg/L | 0.20 |
| Sediment (freshwater) | Local PEC: 15.6 mg/kg | 0.43 |
| Sewage treatment plant | Not relevant (no STP) | |
| Marine water | Not relevant (no emissions to marine water) | |
| Sediment (marine water) | Not relevant (no emissions to marine water) | |
| Agricultural soil | Local PEC: 10.0 mg/kg dw | 0.40 |

Assessment of exposure of man via the environment is not included, since this is for most metal compounds based upon monitoring data rather than modelled data.

9.1.1.4. Guidance to downstream users

If a downstream user (DU) has OCs or RMMs outside the OC/RMM specifications in the generic ES, then the DU can evaluate whether he works inside the boundaries set by the ES through scaling:

- Scaling method, exposure estimation tool used: Fertilizer Environmental Exposure (FEE) Tool v1.2
<http://www.reachfertilizers.com/>
- Scalable parameters:
 - Total annual fertilizer use rate,
 - Number of applications,
 - Time between applications,
 - Crop type,
 - Crop growth stage,
 - European crop yield scenario,
 - Crop substance concentration,
 - Crop yield,
 - Risk management measures (drift and runoff reduction, soil incorporation).All other parameters have to be taken directly from the exposure scenario provided.
- Boundaries of scaling: refer to boundaries as set in Fertilizer Environmental Exposure (FEE) Tool v1.2
<http://www.reachfertilizers.com/>
- For Scaling instructions please go to the following website: <http://www.reachfertilizers.com/>